



## Monroe County Soil & Water Conservation District

1200A Scottsville Rd, Suite 160

Rochester, NY 14624

Ph: 585-473-2120 x 3

[www.monroecountyswcd.org](http://www.monroecountyswcd.org)

# Fish Pond Information

Creating a fish pond can be an easy and rewarding venture. The success of your fish pond relies greatly on the habitat you provide. The type of habitat you create will determine the type of fish that it can support. The following are important characteristics of a fish habitat.

### Temperature

Temperature is fundamental in determining what type of fish will live in your pond. There are two categories of ponds with regards to temperature. Cold water ponds are those with temperatures of 74 degrees or below. Warm water ponds are those with temperatures regularly exceeding 74 degrees. The distinction is made primarily to determine whether a pond can support trout, which require cold water. The temperature of your pond water is influenced by the source of water, geographic location, size and depth. Ponds fed by surface runoff tend to be on the warmer side due to exposure to the sun and warm air before the water enters the pond. Ponds fed by ground water often stay cooler, as groundwater can maintain a much lower temperature in the summer months than surface water. Cold water ponds are almost always fed by some ground water source. Generally, the larger the volume of water present in your pond, the longer it will take to warm in the summer months and the cooler it will stay. In addition, the deeper your pond is the cooler the water will stay.

### Depth

For a warm water pond which supports fish, a minimum depth of 8 feet is necessary over a ½ acre of surface area or greater, a depth of 10 feet is recommended. For a cold water pond that will support trout, 8 to 10 feet of depth over a ½ acre of surface area or greater is recommended for a spring fed pond. 16 feet of depth over a ½ acre or greater is necessary if no spring is present. Don't forget that ice will most probably cover your pond in mid winter. On average in Monroe County, the thickness of ice on a pond will reach 12 inches. If the ice is then covered with snow, sunlight will be prevented from penetrating. If your pond is shallow and therefore contains excessive vegetation, the lack of sunlight will cause the plants to die. As a result, the decomposition of aquatic vegetation will deplete oxygen from the water and your fish will die.

Many species of warm water fish also require shallow waters for spawning. The depth and character of the spawning area is species specific and will be discussed later.

## **Water Quality**

Phosphorous and nitrogen are nutrients which are important to an aquatic ecosystem. Appropriate amounts enrich the growth of aquatic vegetation. The growth of plants provides habitat for forage fish and insect life which are then consumed by predator fish. In addition, vegetation provides spawning areas. Excess nutrients can be detrimental however. Excess nutrients enter your pond through runoff from cropland and livestock holding areas, or from your own fertilized lawn and neighboring residential areas. This can catalyze algae blooms which contribute to the turbidity the water in your pond. As a result, vegetation beneath the surface lacking enough sunlight can die and decompose, thus depleting the water of dissolved oxygen. Dissolved oxygen is a necessary component for fish survival.

A well balanced pond would consist of 1, 2 or 3 game fish or predator species, an adequate forage base, insect life and aquatic vegetation. Fathead Minnows and Golden Shiners can be stocked in combination as forage food or a single species plus additional life such as: frogs, tad poles, crayfish and insect larvae. Fish and other pond life can be ordered through the Monroe County Soil and Water Conservation District office or from local hatcheries. Be sure to request an order form at least 30 days before your stocking date. A farm fish pond license or stocking permit may be required. Contact the New York State Department of Environmental Conservation for further information. (See appendices A and B for contact information and area hatcheries)

## **Warm Water Fish**

**Fish:** Largemouth Bass (*Micropterus salmoides*) Predator Fish

**Preferred Habitat:** Well vegetated areas

**Food Preferences:** insects, frogs, crayfish and small fish

**Spawning Preferences:** Water depth of 1-3 feet and in shallow depressions about 20 inches wide. Spawn in early May to late June when the water temperature reaches 62-64 degrees.

**Stocking Facts:** 100 per surface acre

**Miscellaneous Facts:** Adults reach size of 8-17", females carry 2,000 – 7,000 eggs per pound of body weight, eggs hatch within 7-10 days depending on water temperature

## **Cold Water Fish**

**Fish:** Brook Trout (*Salvelinus fontinalis*) Predator Fish

**Preferred Habitat:** Surface water must stay below 72 degrees in the summer and sustain temperatures below 65 degrees at greater depths, prefer variable underwater habitat (boulders, rock piles, bumps and depressions)

**Food Preferences:** Mollusks, crayfish, other small fish and insects

**Spawning Preferences:** Spawn in the spring in shallow areas of gravel or cobble and require running water

**Stocking Facts:** 300 per surface acre

**Miscellaneous Facts:** Relatively easy to catch and short-lived in relation to Brown and Rainbow Trout, average life expectancy is 3 years

**Fish:** Brown Trout (*Salmo trutta*) Predator Fish

**Preferred Habitat:** Water must stay below 75 degrees in the summer, prefer variable underwater habitat (boulders, rock piles, bumps and depressions)

**Food Preferences:** Mollusks, crayfish, other small fish and insects. Exceptionally large Brown Trout will occasionally feed on frogs, birds and small rodents

**Spawning Preferences:** Spawn in the fall in shallow areas of gravel or cobble, require running water, eggs do not hatch until the spring

**Stocking Facts:** 300 per surface acre

**Miscellaneous Facts:** Very active at night, somewhat difficult to catch during the day

**Fish:** Rainbow Trout (*Oncorhynchus mykiss*) Predator Fish

**Preferred Habitat:** Water temperature must stay below 75 degrees in the summer, can stand long periods at 65-70 degrees but prefer colder.

Variable underwater habitat (boulders, rock piles, bumps and Depressions)

**Food Preferences:** Mollusks, crayfish, other small fish and insects

**Spawning Preferences:** Spawn in the fall in shallow areas of gravel or cobble, require running water, eggs do not hatch until the spring

**Stocking Facts:** 300 per surface acre

**Miscellaneous Facts:** Life expectancy is 3-5 years

### Warm and Cold Water Fish

**Fish:** Yellow Perch (*Perca flavescens*) Predator and Forage Fish

**Preferred Habitat:** Cool, clean water with a sandy or plant covered bottom

**Food Preferences:** Small Perch: zooplankton and insect larvae Larger Perch: crayfish, snails, insects, other small fish and young of its own species.

**Spawning Preferences:** Spawn in shallow, weedy waters at night when the temperature reaches 45-50 degrees in the spring.

**Stocking Facts:** 300 per surface acre

**Miscellaneous Facts:** Average 6-12" in length, females carry 10,000 to 75,000 eggs which are emitted in a gelatinous ribbon over weeds and brush. Due to high fertility, 1/4 to 1/2 of eggs will hatch.

**Fish:** Walleye (*Stizostedion vitreum vitreum*) Predator Fish

**Preferred Habitat:** Requires somewhat cool water with temperatures less than 85 degrees in the summer, they have large light sensitive eyes, so they stay in deeper water during the day and go to the shallows at night.

**Food Preferences:** Young Walleye: zooplankton and insects and their larvae After 2-3 inches in size: small fish of all kinds and crayfish

**Spawning Preferences:** Spawn in April and May when temperatures reach 45-50 degrees, they build nests in colonies in firm sand or fine gravels in water 7 feet deep or less

**Stocking Facts:** 100 per surface acre

**Miscellaneous Facts:** Primarily feed at dusk and dawn, largest member of the Perch family, female carries 25,000-50,000 eggs per pound of body weight, 5-20 percent of the eggs will hatch within 12-18 days depending on water temperature

**Fish:** Fathead Minnow (*Pimephales promelas*) Forage Fish

**Preferred Habitat:** Areas with abundant vegetation

**Food Preferences:** Microscopic algae and other types of plankton

**Spawning Preferences:** Eggs are laid on the underside of boards, rocks or other flat objects

**Stocking Facts:** 2,000 per surface acre

**Miscellaneous Facts:** Valuable forage fish and bait species for all predator fish

**Fish:** Golden Shiner (*Notemigonus crysoleucas*) Forage Fish

**Preferred Habitat:** Clear water with weedy areas

**Food Preferences:** Aquatic insects, mollusks and algae

**Spawning Preferences:** Spawn at numerous intervals in weed beds during midsummer

**Stocking Facts:** 2,000 per surface acre

**Miscellaneous Facts:** Adults grow to a size of 3-7 inches long, considered a valuable forage fish due to the fact that they spawn more than once, producing young throughout the summer

**Crustacean:** Crayfish (*Cambarus cambarus*) Forage Fish

**Preferred Habitat:** Rocky bottoms with places to hide, they are sensitive to light and therefore most active at night.

**Food Preferences:** Small fish and dead crayfish

**Stocking Facts:** 600-1,000 per surface acre

**Miscellaneous Facts:** When a crayfish is molting, their hard shell is discarded. This is when they are most desirable as bait and forage.

Additional information can be found on the Cornell Cooperative Extension Website.

<[www.dnr.cornell.edu/ext/fish/pond1.htm](http://www.dnr.cornell.edu/ext/fish/pond1.htm)>

# Appendix A

## **New York State Department of Environmental Conservation (NYSDEC)**

Region 8 – Division of Regulatory Affairs  
6274 E. Avon-Lima Road  
Avon, NY 14414-9519  
(585) 226-2466  
<[www.dec.state.ny.us](http://www.dec.state.ny.us)>

## **United States Army Corps of Engineers**

Regulatory Branch  
Buffalo District, Corps of Engineers  
1776 Niagara Street  
Buffalo, NY 14207  
(585) 879-4313  
<[www.usace.army.mil](http://www.usace.army.mil)>

## **Monroe County Soil and Water Conservation District**

1200A Scottsville Rd, Suite 160  
Rochester, NY 14624  
(585) 473-2120 x3  
<[www.monroecountyswcd.org](http://www.monroecountyswcd.org)>

## **Cornell Cooperative Extension**

Main Office  
365 Roberts Hall  
Ithaca, New York 14853-5905  
(607) 255-2237  
<[www.cce.cornell.edu](http://www.cce.cornell.edu)>

Monroe County Branch  
249 Highland Ave  
Rochester, NY 14620  
(585) 461-1000

Field Crops Specialist  
(585) 433-2651

## **Monroe County Department of Health**

111 Westfall Road  
Rochester, NY 14620  
(585) 274-6068  
<[www.co.monroe.ny.us/health/](http://www.co.monroe.ny.us/health/)>

## **Dig Safe New York**

1-800-962-7962

# Appendix C

## Fish Hatcheries

Fish can be ordered through the Monroe County Soil and Water Conservation District in early spring, mid-summer and late summer. We provide Triploid Grass Carp, Koi, Golden Shiners, Fathead Minnows and Crayfish. Please call our office at (585) 473-2120 x3 for more information. Fish are also available through the hatcheries listed below. This list is provided for information purposes. The Monroe County Soil and Water Conservation District does not specifically endorse or recommend these businesses.

**Bogner's Hatchery**

2045 Route 5 Utica, NY 13502  
(315)733-3046

Brook Trout

**Finger Lakes Aquaculture**

7627 County Rd 36 Naples, NY 14512  
(585) 374-2974

Bass, Black Crappie, Bluegill, Brook Trout, Rainbow Trout, Fathead Minnows, Golden Shiners, Walleye, Yellow Perch

**Fenton's Smith Creek Fish Farm**

5249 Poplar Tree Rd Bliss, NY 14024  
(585) 322-7703

Bass, Trout, Bluegills, Catfish, Fathead Minnows, Gold Fish, Koi, Perch, Shiners, Walleye, Rosey Red Minnows, Grass Carp

**Empire Fisheries**

4695 Quarry Rd Warsaw, NY 14569  
(585) 786-3315

Large Mouth Bass, Walleye

**George's Bait Farm**

1980 Eastwood Rd East Aurora, NY 14052  
(585) 652-5129

Large Mouth Bass, Fathead Minnows

**Brown's Trout Hatchery**

Route 362 Bliss, NY 14024  
(585) 322-7322

Bass, Trout

**Willow Pond Aqua Farm**

171 Telyea St Canandaigua, NY 14424  
(585) 396-2753

Large Mouth Bass, Trout, Walleye

**Larry Myers**

4825 Federal Rd Hemlock, NY 14466

Large Mouth Bass, Channel Catfish, Yellow Perch, Black Crappie, White Crappie

**Highline Bass**

8050 Garlinghouse Rd Naples, NY 14512  
(585) 374-5114

Large Mouth Bass, Walleye Fingerlings, Trout

**Coolwater Fish Farm**

5636 Lakeville-Groveland Rd Geneseo, NY 14454  
(585) 243-3543

Trout, Walleye, Yellow Perch, Large Mouth Bass, Catfish, Northern Pike, Landlocked Salmon, Fathead Minnows, Golden Shiners, Sunfish

## References

Burkunas, Vytautas, J. Fish Pond Information. Monroe County Soil and Water Conservation District. January, 1993.

Burkunas, Vytautas, J. Personal Interview. May, 2002.

“Farm Fish Pond Management”. New York State Department of Environmental Conservation. 2002. <[dec.state.ny.us/website/dfwmr/fish/farmpdmg.html](http://dec.state.ny.us/website/dfwmr/fish/farmpdmg.html)>

“Fish Pond Management in New York Ponds.” Cornell Cooperative Extension. 2002. <[dnr.cornell.edu/ext/fish/pond1.htm](http://dnr.cornell.edu/ext/fish/pond1.htm)>

“Permits, Applications.” New York State Department of Environmental Conservation. 2002 <[www.dec.state.ny.us/website/dcs/permits\\_level2.html](http://www.dec.state.ny.us/website/dcs/permits_level2.html)>